References

[1] NOAA, 2012. Report on the National Significance of California's Ocean Economy.

[2] Kite-Powell, H, C Colgan, R Weiher, 2008. Estimating the Economic Benefits of Regional Ocean Observing Systems. Coastal Management 36(2):125-145. DOI: 10.1080/08920750701868002

[3] Rayner, R., C Jolly, and C. Gouldman, 2019. Ocean Observing and the Blue Economy. Front. Mar. Sci., https://doi.org/10.3389/fmars.2019.00330

[4] Thoe, W, M Gold, A Griesbach, M Grimmer, ML Taggart, and AB Boehm, 2014. Sunny with a Chance of Gastroenteritis: Predicting Swimmer Risk at California Beaches. Environ. Sci. Technol. 2015, 49, 1, 423–431. https://doi.org/10.1021/es504701j.

[5] Ritzman, J, et al., 2018. Economic and sociocultural impacts of fisheries closures in two fishing-dependent communities following the massive 2015 U.S. West Coast harmful algal bloomHarmful Algae 80, 35-45. https://doi.org/10.1016/j.hal.2018.09.002

[6] Chavez, FP, JT Pennington, RP Michisaki, M Blum, GM Chavez, J Friederich, B Jones, R Herlien, B Kieft, B Hobson, AS Ren, J Ryan, JC Sevadjian, C Wahl, KR Walz, K Yamahara, GE Friederich, and M Messié, 2017. Climate variability and change: Response of a coastal ocean ecosystem. Oceanography 30(4):128–145, https://doi.org/10.5670/oceanog.2017.429.

[7] Bindoff, NL, et al., 2019. Changing Ocean, Marine Ecosystems, and Dependent Communities. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)].

[8] Li, G., Cheng, L., Zhu, J. et al. 2020. Increasing ocean stratification over the past half-century. Nat. Clim. Chang.. https://doi.org/10.1038/s41558-020-00918-2.

[9] Jacox, MG, MA Alexander, NJ Mantua, JD Scott, G Hervieux, RS Webb, and FE Werner, 2018. Forcing of Multiyear Extreme Ocean Temperatures that Impacted California Current Living Marine Resources in 2016. Bull. Amer. Meteor. Soc., 99, S27–S33, https://doi.org/10.1175/BAMS-D-17-0119.1

[10] Jacox, M.G., Alexander, M.A., Bograd, S.J. et al. 2020. Thermal displacement by marine heatwaves. Nature 584, 82–86 https://doi.org/10.1038/s41586-020-2534-z.

[11] Sanford, E, JL Sones, M García-Reyes, JHR Goddard, and JL Largier, 2019. Widespread shifts in the coastal biota of northern California during the 2014–2016 marine heatwaves. Scientific Reports, doi.org/10.1038/s41598-019-40784-3.

[12] Harvey, C, et al., 2020. Ecosystem Status Report of the California Current for 2019-20: A Summary of Ecosystem Indicators Compiled by the California Current Integrated Ecosystem Assessment Team (CCIEA). doi: https://doi.org/10.25923/e5rb-9f55.

[13] Rogers-Bennett, L, CA Catton, 2019. Marine heat waves and multiple stressors tip bull kelp forest to sea urchin barrens. Sci Rep 9, 15050 https://doi.org/10.1038/s41598-019-51114-y

[14] Ryan, JP, RM Kudela, JM Birch, M Blum, HA Bowers, FP Chavez, GJ Doucette, K Hayashi, R Marin III, CM Mikulski, JT Pennington, CA Scholin, GJ Smith, A Woods, Y Zhang, 2017. Causality of an extreme harmful algal bloom in Monterey Bay, California, during the 2014–2016 northeast Pacific warm anomaly, Geophys. Res. Lett., 44, doi:10.1002/2017GL072637.

[15] Anderson, CR, E Berdalet, R Kudela, CK Cusack, J Silke, E O'Rourke, D Dugan, M McCammon, JA Newton, SK Moore, and K Paige, 2019. Scaling up from regional case studies to a global harmful algal bloom observing system. Frontiers in Marine Science, 6, p250.

[16] Fischer, AD, K Hayashi, A McGaraghan, and R Kudela, 2020. Return of the "age of dinoflagellates" in Monterey Bay: Drivers of dinoflagellate dominance examined using automated imaging flow cytometry and long-term time series analysis. Limnology and Oceanography 9999, 1-17.

[17] McCabe, RM et al. 2016 An unprecedented coastwide toxic algal bloom linked to anomalous ocean conditions. Geophysical Research Letters. https://doi.org/10.1002/2016GL070023.

[18] Kudela, RM, J. Lane, and WP Cochlan, 2008. The potential role of anthropogenically derived nitrogen in the growth of harmful algae in California, USA. Harmful Algae 8 (1):103-110. doi: https://doi.org/10.1016/j.hal.2008.08.019.

[19] Feely, RA, CL Sabine, J Martin Hernandez-Ayon, D Ianson, and B Hales, 2008. Evidence for Upwelling of Corrosive & Acidified Water onto the Continental Shelf. Science 320 (5882):1490. doi: 10.1126/science.1155676.

[20] Chan, F, et al, 2017. Persistent spatial structuring of coastal ocean acidification in the California Current System. Sci Rep, 7(1), 2526. doi:10.1038/s41598-017-02777-y.

[21] Breitburg, D, et al. 2018. Declining oxygen in the global ocean and coastal waters. Science 05 359, DOI: 10.1126/science.aam7240.

[22] Weisberg, SB, F Chan, JP Barry, AB Boehm, S Busch, SR Cooley, RA Feely, LA Levin, 2020. Enhancing California's Ocean Acidification and Hypoxia Monitoring Network. California Ocean Science Trust, Sacramento, California, USA.

[23] Osborne EB, RC Thunell, N Gruber, RA Feely, and CR Benitez-Nelson, 2020. Decadal variability in twentieth-century ocean acidification in the California Current Ecosystem. Nature Geoscience 18: 43–49.

[24] Reynolds, CA, JD Doyle, FM Ralph, R Demirdjian, 2019. Adjoint Sensitivity of North Pacific Atmospheric River Forecasts., Mon. Wea. Rev. 147 (6): 1871–1897. https://doi.org/10.1175/MWR-D-18-0347.1

[25] Chao, Y, et al., 2017. Development, implementation, and validation of a California coastal ocean modeling, data assimilation, and forecasting system Deep Sea Research Part II: 151: 49-63. https://doi.org/10.1016/j.dsr2.2017.04.013.

[26] Moore, AM, CA Edwards, J Fiechter, P Drake, HG Arango, E Neveu, S Gurol and AT Weaver. (2013) A 4D-Var Analysis System for the California Current: A Prototype for an Operational Regional Ocean Data Assimilation System. In ``Data Assimilation for Atmospheric, Oceanic and Hydrological Applications, Vol. II," Liang Xu and Seon Park, Eds. Springer, Chapter 14, 345-366.

[27] Iwamoto, MM, J Dorton, J.Newton, M Yerta, J Gibeaut, T Shyka, B Kirkpatrick, and R Currier, 2019. Meeting regional, coastal and ocean user needs with tailored data products: A stakeholder-driven process. Frontiers in Marine Science 6:290. https://doi.org/10.3389/fmars.2019.00290.

[28] IOOS, 2018. US IOOS Enterprise Strategic Plan (2018-2022), NOAA, Silver Spring MD.

[29] Roarty, H, T Cook, L Hazard, D George, J Harlan, S Cosoli, et al., 2019. The global high frequency radar network. Front. Mar. Sci. 6:164. doi: 10.3389/fmars.2019.00164.

[30] Testor P, et al., 2019. OceanGliders: A Component of the Integrated GOOS. Front. Mar. Sci. 6:422. doi: 10.3389/fmars.2019.00422.

[31] NASA, 2020. https://appliedsciences.nasa.gov/what-we-do/ecological-forecasting

[32] Levin, LA, Bette, BJ et al. 2019. Global Observing Needs in the Deep Ocean. Frontiers in Marine Science 6 (241). doi: 10.3389/fmars.2019.00241.

[33] Tilbrook, B, Jewett, EB et al. 2019. An Enhanced Ocean Acidification Observing Network: From People to Technology to Data Synthesis and Information Exchange. Frontiers in Marine Science 6 (337). doi: 10.3389/fmars.2019.00337.

[34] Muller-Karger, FE et al., 2018. Advancing Marine Biological Observations and Data Requirements of the Complementary Essential Ocean Variables (EOVs) and Essential Biodiversity Variables (EBVs) Frameworks. Front. Mar. Sci., 27, https://doi.org/10.3389/fmars.2018.00211.

[35] Harcourt, R, et al., 2019. Animal-Borne Telemetry: An Integral Component of the Ocean Observing Toolkit, Frontiers in Marine Science, 10.3389/fmars.2019.00326, 6, .

[36] Jacox, MG, CA Edwards, EL Hazen, and SJ Bograd, 2018. Coastal upwelling revisited: Ekman, Bakun, and improved upwelling indices for the U.S. West Coast. Journal of Geophysical Research: Oceans, 123, 7332-7350. https://doi.org/10.1029/2018JC014187.

[37] National Agricultural Statistics Service. 2014. "2013 Census of Aquaculture." *United States Department of Agriculture*. Vol. 3. https://www.agcensus.usda.gov/Publications/2012/Online_Resources/Aquaculture/Aqua.pdf

[38] Anderson, CR, RM Kudela, et al., 2016. Initial skill assessment of the California harmful algae risk mapping (C-HARM) system. Harmful Algae, 59, pp.1-18.

[39] Ryan, JP, et al., 2019. Integrative passive acoustic monitoring in Monterey Bay National Marine Sanctuary. The Journal of the Acoustical Society of America 146, 2772; https://doi.org/10.1121/1.5136603.

[40] Ocean Science Trust, 2016. Frequently Asked Questions: Harmful Algal Blooms and California Fisheries, Developed in Response to the 2015-2016 Domoic Acid Event. California Ocean Science Trust, Oakland, CA.

[41] Barton, A, GG Waldbusser, RA Feely, SB Weisberg, JA Newton, B Hales, S Cudd, B Eudeline, CJ Langdon, I Jefferds, T King, A Suhrbier, K McLaughlin, 2015. Impacts of coastal acidification on the Pacific Northwest shellfish industry and adaptation strategies implemented in response. Oceanography, 28(2), 146-159. doi:10.5670/oceanog.2015.38.

[42] Swezey, DS, et al., 2020. Evolved differences in energy metabolism and growth dictate the impacts of ocean acidification on abalone aquaculture. Proceedings of the National Academy of Sciences, 117 (42), 26513-26519. https://doi.org/10.1073/pnas.2006910117.

[43] Sievanen, L, et al., 2018. California's Coast and Ocean Summary Report. California's Fourth Climate Change Assessment. Publication number: SUMCCC4A-2018- 011.

[44] Hazen, EL, et al., 2019. Marine top predators as climate and ecosystem sentinels. Front. Ecol. Environ. 17, 565–574.

[45] Keates, Theresa R., Raphael M. Kudela, Rachel R. Holser, Luis A. Hückstädt, Samantha E. Simmons, and Daniel P. Costa. 2020. "Chlorophyll fluorescence as measured in situ by animal-borne instruments in the northeastern Pacific Ocean." Journal of Marine Systems 203:103265. doi: https://doi.org/10.1016/j.jmarsys.2019.103265.

[46] Weisberg, SB, F Chan, JP Barry, AB Boehm, S Busch, SR Cooley, RA Feely, LA Levin, 2020. Enhancing California's Ocean Acidification and Hypoxia Monitoring Network. California Ocean Science Trust, Sacramento, California, USA.

[47] NOAA, 2020. U.S. West Coast SCCOOS • CeNCOOS • NANOOS - ATN • MBON • OTN Biological Observations Workshop Summary Report: Identifying Regional Needs and Priorities for Animal Telemetry and Biodiversity Observations of Aquatic Species, Silver Spring, MD.

[48] Kishi, MJ, et al., 2007. NEMURO - A lower trophic level model for the north pacific marine ecosystem. Ecol. Modell. 202, 12-25. doi:10.1016/j.ecolmodel.2006.08.021.

[49] Manzer, CR, TP Connolly, E McPhee-Shaw, GJ Smith, 2019. Physical factors influencing phytoplankton abundance in southern Monterey Bay. Continental Shelf Res.180:1-13. DOI: 10.1016/j.csr.2019.04.007

[50] Friederich, GE, PG Brewer, R Herlien, FP Chavez, 1995. Measurement of sea surface partial pressure of CO2 from a moored buoy. Deep Sea Res. Part I, 42: 1175-1186, 10.1016/0967-0637(95)00044-7.

[51] Jahncke, J, BL Saenz, CL Abraham, C Rintoul, RW Bradley, WJ Sydeman, 2008. Ecosystem responses to short-term climate variability in the Gulf of the Farallones, California. Prog Oceanogr. 77: 182-193.

[52] Trainer, VL, RM Kudela, MV Hunter, NG Adams and RM Mccabe (in press) Climate extreme seeds a new domoic acid hotspot on the US west coast. Front. Clim., doi: 10.3389/fclim.2020.571836.

[53] Kudela, RM, A Bickel, ML Carter, MDA Howard, L Rosenfeld, 2015. The Monitoring of Harmful Algal Blooms through Ocean Observing: the Development of the California Harmful Algal Bloom Monitoring and Alert Program. Invited chapter to Coastal Ocean Observing Systems, Y Liu, H Kerkering, RH Weisberg (eds.). Elsevier, Chapter 5, pp. 58-75. ISBN 978-0-12-802022-7.

[54] Ocean Science Trust, 2016. Framing the Scientific Opportunities on Harmful Algal Blooms and California Fisheries Scientific Insights, Recommendations and Guidance for California. California Ocean Science Trust, Oakland, CA.

[55] Mazzini, PLF, JA Barth, RK Shearman, and A Erofeev, 2014. Buoyancy-driven coastal currents off Oregon during fall and winter. J. Phys. Oceanogr. 44, 2854–2876. doi: 10.1175/JPO-D-14-0012.1.

[56] Rudnick, DL, RE Davis, and JT Sherman, 2016. Spray underwater glider operations. J. Atmospheric Oceanic Technol. 33, 1113–1122. doi: 10.1175/JTECH-D-15-0252.1.

[57] Takeshita, Y et al., (in press) Accurate pH and O2 measurements from Spray underwater gliders. Journal of Atmospheric and Oceanic Technology.

[58] Bittig HC, et al., 2019. A BGC-Argo Guide: Planning, Deployment, Data Handling and Usage. Front. Mar. Sci. 6:502. doi: 10.3389/fmars.2019.00502

[59] Ohman, M.D., Davis, R.E., Sherman, J.T., Grindley, K.R., Whitmore, B.M., Nickels, C.F. and Ellen, J.S. (2019), *Zooglider:* An autonomous vehicle for optical and acoustic sensing of zooplankton. Limnol Oceanogr Methods, 17: 69-86. https://doi.org/10.1002/lom3.10301

[60] Chapple, TK, 2011. A first estimate of white shark, *Carcharodon carcharias*, abundance off Central California. Biol. Lett.: 7: 581-583.

[61] CDFW, 2020, California Department of Fish And Wildlife: White Shark Information. https://wildlife.ca.gov/Conservation/Marine/White-Shark.

[62] NOAA, 2020. NOAA Unmanned Systems Strategy: Maximizing Value for Science-based Mission Support, Silver Spring, MD.

[63] Anruszkiewicz, EA, HA Starks, FP Chavez, LM Sassoubre, BA Block, AB Boehm, 2017. Biomonitoring of marine vertebrates in Monterey Bay using eDNA metabarcoding. PLoS ONE 12(4): e0176343. https://doi.org/10.1371/journal.pone.0176343.

[64] Ausubel, JH, MY Stoeckle, PG Gaffney II, 2019. Final Report, First National Conference on Marine Environmental DNA. https://phe.rockefeller.edu/research/e-dna/.

[65] Djurhuus, A, et al. 2020. Environmental DNA reveals seasonal shifts and potential interactions in a marine community. Nature Comms.: 11:254: https://doi.org/10.1038/s41467-019-14105-1.

[66] NOAA, 2020. NOAA 'Omics Strategy Strategic Application of Transformational Tools, Silver Spring, MD.

[67] Lombard F, et al., 2019. Globally Consistent Quantitative Observations of Planktonic Ecosystems. Front. Mar. Sci. 6:196. doi: 10.3389/fmars.2019.00196.

[68] Lamouroux, J, G Charria, P De Mey, S Raynaud, C Heyraud, P Craneguy, et al., 2016. Objective assessment of the contribution of the RECOPESCA network to the monitoring of 3D coastal ocean variables in the Bay of Biscay and the English Channel. *Ocean Dyn.* 66, 567–588. doi: 10.1007/s10236-016-0938-y.

[69] Charria, G, S Theetten, F Vandermeirsch, Ö Yelekçi, and N Audiffren, 2017. Interannual evolution of (sub)mesoscale dynamics in the Bay of Biscay. *Ocean Sci.*13, 777–797. doi: 10.5194/os-13-777-2017.

[70] NOAA, 2020. NOAA Cloud Strategy: Maximizing the Value of NOAA's Cloud Services, Silver Spring, MD.

[71] IOOS, 2020, Data Standards & Requirements (https://ioos.noaa.gov/data/contribute-data/), NOAA, Silver Spring MD.

[72] NOAA, 2020. NOAA Artificial Intelligence Strategy: Analytics for Next Generation Earth Science, Silver Spring, MD.

[73] Chao, Y, JD Farrara, H Zhang, YJ Zhang, E Ateljevich, F Chai, CO Davis, R Dugdale, F Wilkerson, 2017. Development, implementation, and validation of a modeling system for the San Francisco Bay and Estuary. Estuarine, Coastal and Shelf Science, 194, 40-56, 2017.

[74] Chao, Y, JD Farrara, E Bjorkstedt, F Chai, F Chavez, DL Rudnick, W Enright, JL Fisher, WT Peterson, GF Welch, CO Davis, RC Dugdale, FP Wilkerson, H Zhang, Y Zhang, E Ateljevich, 2018. The origins of the anomalous warming in the California coastal, ocean and San Francisco Bay during 2014–2016, J. Geophys. Res. Oceans, 122, doi:10.1002/2017JC013120.

[75] NOAA, 2020f San Francisco Bay Operational Forecast System (SFBOFS): https://tidesandcurrents.noaa.gov/ofs/sfbofs/sfbofs.html. Center for Operational Oceanographic Products and Services, Silver Spring, MD.

[76] Dugdale, RC, FP Wilkerson, AE Parker, 2013. A biogeochemical model of phytoplankton productivity in an urban estuary: the importance of ammonium and freshwater flow. Ecological Modelling, 263, 291-307.

[77] Wang, Z, F Chai, R Dugdale, et al., 2020. The interannual variabilities of chlorophyll and nutrients in San Francisco Bay: a modeling study. Ocean Dynamics 70, 1169–1186. https://doi.org/10.1007/s10236-020-01386-0.

[78] NOAA, NMFS. 2020. Large Whale Entanglements off the U.S. West Coast, From 1982–2017. NOAA Technical Memorandum NMFS-OPR-63

[79] Rockwood RC, J Calambokidis, J Jahncke, 2017. High mortality of blue, humpback and fin whales from modeling of vessel collisions on the U. S. West Coast suggests population impacts and insufficient protection. PLoS ONE 12(8): e0183052. https://doi.org/10.1371/journal.pone.0183052

[80] Northern Economics, Inc., 2013. The Economic Impact of Shellfish Aquaculture in Washington, Oregon and California. Prepared for Pacific Shellfish Institute.

[81] Sydeman, WJ, SA Thompson, M García-Reyes, M Kahru, WT Peterson, and JL Largier, 2014. Multivariate ocean-climate indicators (MOCI) for the central California Current: Environmental change, 1990–2010. Progress in Oceanography 120: 352-369.

[82] García-Reyes, M, and WJ Sydeman, 2017. California multivariate ocean climate indicator (MOCI) and marine ecosystem dynamics. Ecological Indicators, 72, pp.521-529.

[83] Hazen, EL, DM Palacios, KA Forney, EA Howell, E Becker, AL Hoover, L Irvine, M DeAngelis, SJ Bograd, BR Mate, and H Bailey, 2017. WhaleWatch: a dynamic management tool for predicting blue whale density in the California Current. J Appl Ecol, 54: 1415-1428. https://doi.org/10.1111/1365-2664.12820

[84] Bell, TW, JA Allen, KC Cavanaugh, DA Siegel, 2020. Three decades of variability in California's giant kelp forests from the Landsat satellites. Remote Sensing of Environment. 238, 110811.

[85] Hamilton, S, TW Bell, J Watson, K Grorud-Colvert, B Menge, 2020. Remote Sensing: Generation of Long-Term Kelp Forest Datasets for Evaluation of Impacts of Climatic Variation. Ecology, 101, e03031.

[86] Welch H, Hazen EL, Bograd SJ, et al., 2019. Practical considerations for operationalizing dynamic management tools. J Appl Ecol.;56:459–469. https://doi.org/10.1111/1365-2664.13281.